

MGI Michiana Gastroenterology Inc.

Telephone: (574) 234-0049

Fax: (574) 234-0053

www.michianagastro.com

Colonoscopy Instructions with Movi Prep

****Please be aware, there could be multiple separate charges for this procedure. One charge for the Facility and one for the Doctor. Possible charges for anesthesia and pathology.****

Patient: _____ Dr. _____

Date: ____ / ____ / ____

Arrival Time: _____ am/pm

Procedure Time: _____ am/pm

____ Michiana Endoscopy Center
53830 Generations Drive
South Bend, IN 46635

____ Plymouth-SJRM C
Register at Surgery Ctr.
Building 5

____ Memorial Hospital
Main Entrance
Patient Registration

____ Mishawaka - SJRMC
Main Entrance
Patient Registration

____ LaPorte Surgical Center
900 I Street
LaPorte, IN 46350

Do these things 7 days before your procedure:

- Stop iron medication and iron supplements.
- If you take Effient, check with the doctor who ordered it to see if you can stop taking it 7 days prior to the procedure.

Do these things 4 days before your procedure:

- Arrange to have someone drive you home. You will receive medicine to keep you relaxed during your procedure, so you cannot drive home or take a cab home alone.
- Stop taking aspirin. It is okay to take Tylenol (Acetaminophen).
- If you take Coumadin or any other blood thinner other than Plavix, check with the doctor who ordered it to see if you can stop taking it for the 4 days before the procedure.
- If you take insulin call the doctor who ordered it for any adjustments you need to make the day of prep and the procedure day. We do not want your blood sugar to be too low.

Do these things 1 day before your procedure (Prep Day):

- DO NOT EAT ANY SOLID FOOD TODAY**
- You may only have clear liquids all day and night. **NO RED OR PURPLE**. Clear liquids are: tea or coffee (no milk or creamer), broth, apple juice, white cranberry juice, 7 up, ginger ale and water. You may also have Gatorade, popsicles and Jell-O, just remember no red or purple.
- Do not drink alcohol.
- Start drinking your prep according to your doctor's instructions, which are located on the back of this page.
 - Note: Some people say taking sips of broth or sucking on a lime in between glasses of prep help to clear the taste.
 - If you feel too full, drink the prep every 30 minutes.
- Keep drinking clear liquids for the rest of the day. This will help the prep work better for you.
- The prep you are drinking will make you go to the bathroom many times. This is needed so your colon will be cleaned out for the procedure.
- You may use Vaseline, any diaper rash cream or baby wipes, etc...on your bottom. This will help it not get so sore.

Do these things on the day of your procedure:


- DO NOT EAT ANY SOLID FOOD**
- You may continue drinking clear liquids, but **stop drinking 3 hours before your procedure**.
- Take your usual morning pills except for Coumadin and Aspirin.
- DO NOT take your diabetic pill(s).
- If you usually check your blood sugar, check it this morning.
- Bring the person who is driving you home with you to the Endoscopy Center.
- Your arrival time to your discharge time is about 2 1/2 hours.
- Bring insurance card, photo id and all medications to your appointment.

OVER ➔

Day Before Procedure
Follow printed instructions on this sheet not the colon prep packaging

1. **Do not eat any solid food today.** It is important to drink at least eight (8) glasses of clear liquids throughout the day.
2. In the morning,


- Empty 1 Pouch A and 1 Pouch B into the disposable container
- Add lukewarm drinking water to the top line of the container. Mix to dissolve



Put the container in the refrigerator to chill.

3. At 6:00 pm, begin drinking the Moviprep as listed below. Be sure to drink additional fluids afterwards.

- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is complete
- Drink 16 oz of the clear liquid of your choice




4. The prep could take effect within 20 minutes, so stay close to the bathroom.
5. Mix second liter of Moviprep after the first liter is completely finished. Use the supplied container again empty 1 pouch A and 1 Pouch B into the disposable container. Add water to the top line of the container and mix to dissolve. Refrigerate for later.

Day of Procedure

1. If you awaken with solid stool, please call (574) 271-0893.
2. **DO NOT EAT ANY SOLID FOOD until after your procedure.**
3. Drink second liter of Moviprep at _____ AM as listed below. Be sure to drink additional fluids afterwards.

- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is complete
- Drink 16 oz of the clear liquid of your choice



4. Drinking clear liquids is encouraged to prevent dehydration.
5. **DO NOT DRINK 3 HOURS BEFORE YOUR PROCEDURE!!**

If you need to cancel, reschedule or have questions regarding the prep, please call the office at 574-234-0049.

IN CASE OF AN EMERGENCY, GO TO THE HOSPITAL EMERGENCY ROOM!